

AD-A274 695 PAGE

Form Approved  
OMB No. 0704-0188

2

Public reporting  
burden for this  
collection of information  
is estimated to average  
1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, reviewing and collecting the data, and reviewing and collecting the data.  
Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington, DC 20503, Paperwork Reduction Project (0704-0188).per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, reviewing and collecting the data, and reviewing and collecting the data.  
Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington, DC 20503, Paperwork Reduction Project (0704-0188).

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE January 8, 1994		3. REPORT TYPE AND DATES COVERED Technical Report	
4. TITLE AND SUBTITLE Transition Metals in Highly Correlated Discotic Phases: Designing Metallomesogens With Selected Intermolecular Organizations				5. FUNDING NUMBERS N00014-92-J-1416	
6. AUTHOR(S) Hanxing Zheng, Chung K. Lai, and Timothy M. Swager				R&T Code 400X098YIP01	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry University of Pennsylvania Philadelphia, PA 19104-6323				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Dr. H. Guard (Program Director) Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000				10. SPONSORING/MONITORING AGENCY REPORT NUMBER Technical Report No. 4	
11. SUPPLEMENTARY NOTES <u>Chemistry of Materials</u> in press.					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. ABSTRACT (Maximum 200 words) Diketonate Schiff-base complexes were investigated, and the square planar derivatives (M=Ni, Cu, Pd) were found to display discotic hexagonal disordered phases ( $D_{hd}$ ). The M=VO analogs are not liquid crystalline. None of the complexes can exhibit a disc-shape independently, and in the mesophase they must organize into a superstructure which on time average leads to antiparallel alignment of nearest neighbors. These discotic antiphases exhibit a number of important properties including low melting points, low viscosity, and close intermolecular contacts of 3.6Å. Additionally when put between two untreated slides, these mesophases were found to readily align with the column axes perpendicular to the surfaces. The Cu and Ni complexes exhibited very similar mesophases and lattice constants whereas the Pd complexes displayed larger intercolumnar distances and higher clearing temperatures. The differences are ascribed to greater intermolecular associations displayed by the Pd complexes. X-ray diffraction studies revealed the Ni and the Cu complexes to have an additional halo centered at $\sim 7.2\text{\AA}$ which may be the result of a doubling of the period along the column axes resulting from the antiparallel arrangement of the antiphase superstructure.					
14. SUBJECT TERMS				15. NUMBER OF PAGES	
				16. PRICE CODE NA	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified		19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	
				20. LIMITATION OF ABSTRACT Unlimited	

DTIC  
ELECTE  
JAN 21 1994  
S E D

94-01800



3P8

94 1 19 004

**OFFICE OF NAVAL RESEARCH**

Grant N00014-92-J-1416

R&T Code 400X098YIP01

Technical Report No. 4

**Transition Metals in Highly Correlated Discotic Phases: Designing  
Metallomesogens With Selected Intermolecular Organizations**

by

Hanxing Zheng, Chung K. Lai, and Timothy M. Swager

Chemistry of Materials in press

Department of Chemistry  
University of Pennsylvania  
Philadelphia, PA 19104-6323

January 8, 1994

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification .....	
By .....	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

**DTIC QUALITY INSPECTED 1**

# Transition Metals in Highly Correlated Discotic Phases: Designing Metallomesogens With Selected Intermolecular Organizations

Hanxing Zheng, Chung K. Lai, and Timothy M. Swager\*

Department of Chemistry  
University of Pennsylvania  
Philadelphia, PA 19104-6323

## Abstract:

The diketone Schiff-base complexes shown below were investigated, and the square planar derivatives ( $M=\text{Ni}$ ,  $\text{Cu}$ ,  $\text{Pd}$ ) were found to display discotic hexagonal disordered phases ( $D_{hd}$ ). The  $M=\text{VO}$  analogs are not liquid crystalline. None of the complexes can exhibit a disc-shape independently, and in the mesophase they must organize into a superstructure which on time average leads to antiparallel alignment of nearest neighbors. These discotic antiphases exhibit a number of important properties including low melting points, low viscosity, and close intermolecular contacts of  $3.6\text{\AA}$ . Additionally when put between two untreated slides, these mesophases were found to readily align with the column axes perpendicular to the surfaces. The  $\text{Cu}$  and  $\text{Ni}$  complexes exhibited very similar mesophases and lattice constants whereas the  $\text{Pd}$  complexes displayed larger intercolumnar distances and higher clearing temperatures. The differences are ascribed to greater intermolecular associations displayed by the  $\text{Pd}$  complexes. X-ray diffraction studies revealed the  $\text{Ni}$  and the  $\text{Cu}$  complexes to have an additional halo centered at  $\sim 7.2\text{\AA}$  which may be the result of a doubling of the period along the column axes resulting from the antiparallel arrangement of the antiphase superstructure.

